

§18. Study of Effect of Plasma Shape and MHD Equilibrium Limit

Suzuki, Y., Yamada, H., Watanabe, K.Y.,
Sakakibara, S., Nakajima, N.,
Yamada, G., Nakamura, Y. (Kyoto Univ.)

Plasma shape control is a major knob to investigate confinement as well as MHD characteristics of magnetically confined plasmas. In particular, plasma elongation has been widely explored in many experiments and theoretical investigations. For example, optimization in the line of W7-AS and W7-X owes much to reduction of Pfirsch-Schluter currents due to elongation. It is also widely recognized that plasma elongation affects energy confinement in tokamaks, which can be seen in the ITER H-mode scaling.

In heliotron plasmas, when the shape of flux surfaces is elongated vertically (horizontally), following characteristic are predicted by numerical and analytical analyses based on the stellarators expansions: (i) the rotational transform on the axis is decreased, (ii) the Pfirsch-Schluter current is suppressed (enhanced), (iii) the Shafranov shift toward the outside of the torus is small (large). Since the Shafranov shift is enhanced by horizontal elongation, equilibrium β limit is lowered in oblate configuration. In general, the equilibrium β limit is defined by the Shafranov shift as large as a half the minor radius, when the deformation of magnetic flux surface is thought not to be tolerable. However, this is

not physically reasonable definition. In the tokamak case, the separatrix is generated inboard side due to enhanced Pfirsch-Schluter currents. Oblate configuration provides the experimental evidence what happens by a large Shafranov shift and can mitigate the approach to the question what the equilibrium β limit is.

In order to consider equilibrium β limit, we study an oblate configuration ($B_Q=150\%$). This configuration is elongated by the external quadrupole field. If the horizontal elongation is superimposed strongly, vacuum magnetic axes were split. Those configurations had already studied in the theoretical and experimental analyses¹⁾²⁾. In this study, since we are interesting the marginal region to equilibrium β limit, we adapt this configuration. Figure 1 shows time evolutions of the electron temperature and density profiles in the typical shot (#68866). The diamagnetic beta $\langle\beta\rangle_{\text{dia}}$ achieved to about 2.5% at $t\sim 1.1$ [s]. The axis on the horizontal cross section shifts $R\sim 4.1$ m. Special notices are the fluttering of the electron temperature profile at $t=1.921$ [s]. This suggests a possibility of appearing of the separatorix in core plasma and equilibrium β limit.

1) Suzuki, Y., et al., Nucl. Fusion **46**, (2006) 123

2) Yamada, H., et al., Fusion Sci. Technol **51**, (2007) 138




Fig. 1. Time evolutions of profiles of the electron temperature and density on the horizontal cross section at highest- β discharge. At $t=1.921$ [s], the fluttering of T_e profile was appeared.